arXiv:1305.0131v1 [cond-mat.mes-hall] 1 May 2013

Classification of engineered topological superconductors

Panagiotis Kotetes*
Institut für Theoretische Festkörperphysik, Karlsruhe Institute of Technology, 76128 Karlsruhe, Germany

I perform a complete classification of 2d, quasi-1d and 1d topological superconductors which originate from the suitable combination of inhomogeneous Rashba spin-orbit coupling, magnetism and superconductivity. My analysis reveals alternative types of topological superconducting platforms for which Majorana fermions are accessible. Specifically, I observe that for quasi-1d systems with Rashba spin-orbit coupling and time-reversal violating superconductivity, Majorana fermions can in priniciple emerge *even in the absence* of magnetism. Furthermore, for the classification I also consider situations where additional "hidden" symmetries emerge, with a significant impact on the topological properties of the system. The latter, generally originate from a combination of space group and complex conjugation operations that separately do not leave the Hamiltonian invariant. Finally, I suggest alternative directions in topological quantum computing for systems with additional unitary symmetries.

PACS numbers: 74.78.-w, 74.45.+c, 03.67.Lx

Introduction

The breakthrough concept of emergent Majorana fermions (MFs) in artificial topological superconducting (TSC) devices, pioneered by Fu and Kane [1], motivated a number of recent experiments [2] that have already provided the first promising results. The two authors demonstrated that the helical surface states of a three-dimensional topological insulator [3], with proximity induced superconducting gap Δ, behave as a time-reversal ($\mathcal{T}$) invariant TSC. When a magnetic field is applied perpendicular to the topological surface, a single MF appears per superconducting vortex. Shortly after this proposal, it was recognized that the catalytic presence of spin-momentum locking could be alternatively provided by the spin-orbit interaction of a Rashba semiconductor [4, 5]. In the case of a semiconducting wire [5], fabricated for instance from InSb, a Zeeman energy $\mu_s|\boldsymbol{\mathcal{B}}|$ is sufficient to lead to MFs localized at the edges. The existence of confined and protected edge MFs is crucial for applications in topological quantum computing [6, 7]. A pair of MFs defines a topological qubit, which is in principle free from decoherence and protected against noise, in stark contrast to traditional spin [8] and superconducting qubits [9]. Furthermore, edge MFs can also give rise to unique transport signatures [10–13], such as the usual [10, 11] or the magnetically controlled [12] 4π-Josephson effect.

In the case of a semiconducting wire [5] with proximity induced superconductivity, the system transits to the topological phase when the criterion $\mu_s|\boldsymbol{\mathcal{B}}| > \sqrt{\mu^2+|\Delta|^2}$ is satisfied (μ defines the chemical potential). The concomitant requirement of a high magnetic field can impede the nanofabrication of the device or restrict the possible manipulations on the MFs. In fact, several proposals concerning quantum information processes rely on the application of strong antiparallel magnetic fields at the nanoscale level [14], something not easily realizable in the lab. As an answer to these obstacles alternative types of engineered TSCs have been put forward, which support MFs without the necessary presence of spin-orbit coupling or the application of a magnetic field [15–19]. In most of these proposals, some kind of inhomogeneous magnetic order coexists with intrinsic or proximity induced superconductivity. For some of these models [18, 19] it has been argued that there is a mapping to the case of the semiconducting wire-based TSC mentioned above. Nonetheless, this mapping does not necessarily imply an exact topological equivalence. In order to discern if two topological systems are equivalent, the topological class and the role of dimensionality must be simultaneously taken into consideration.

In this manuscript I present a complete topological classification of low-dimensional TSCs that support MFs and originate from the combined presence of inhomogeneous Rashba spin-orbit coupling $v(\boldsymbol{r})$, magnetism $\boldsymbol{M}(\boldsymbol{r})$ and superconductivity $\Delta(\boldsymbol{r})$. My primary goal is to shed light on the topological connection between different existing proposals for engineered TSCs and in addition to propose alternative advantageous platforms. For my analysis I will consider 2d, quasi-1d and 1d systems. The quasi-1d case is obtained from the strict 2d case by the inclusion of a confining potential $V(\boldsymbol{r})$. My study provides new engineered TSC that are experimentally accessible. For example, I find that in the case of a quasi-1d system with Rashba spin-orbit coupling and time-reversal symmetry ($\mathcal{T}$) violating

---

*Electronic address: panagiotis.kotetes@kit.edu

superconducting order parameter, MFs can emerge even in the absence of magnetic fields or any type of inhomogeneous magnetism. In addition, I examine the effects of dimensionality on the robustness of MFs through separating the systems under investigation into weak and strong engineered TSCs. Furthermore, I illustrate that so far overlooked discrete symmetries, that I shall refer to as "hidden" symetries ($\mathcal{O}$), distinguish models previously considered as topologically equivalent. Generally, hidden symmetries can be either unitary or anti-unitary and result from a combination of space group, time-reversal or other internal symmetry operations that when considered separately do not leave the Hamiltonian invariant (e.g. [20, 21]). Here I discuss two examples of hidden symmetries: i) a unitary hidden symmetry resulting from the combination of a reflection and a translation and ii) an anti-unitary symmetry resulting from the combination of time-reversal and translation operations. Finally, I also discuss new topological quantum computing (TQC) perspectives that appear when hidden symmetries are present.

At this point, I give a brief description of how the several sections are organized. In Section I, I provide a short introduction to Majorana fermions and introduce the general Hamiltonian that describes the systems of interest. In Section II, I shortly review the topological classification methods with special focus on the situations where additional unitary and anti-unitary symmetries are present. In Section III, I present an overview of my main results (Table II) concerning the classification of TSCs when all possible spatial symmetries are broken. I further discuss how the emergence of hidden symmetries can modify Table II. In Section IV, I provide a detailed analysis and justification of the results presented in Section III. In Section V, I present two specific examples of systems characterized by a hidden symmetry and study the impact of the latter on the topological properties. In Section VI, I discuss how the presence of hidden symmetries can be useful for developing topological quantum computing protocols and suggest possible candidate systems that could be used for this purpose. Finally, Section VII summarizes my main results and related conclusions.

I. MAJORANA FERMIONS AND MODEL HAMILTONIAN

In condensed matter physics MFs are not fundamental particles [22] but excitations of a many-body system [23, 24]. Essentially, what we define as MFs are the operators γ_α (α is just a label) which satisfy $\{\gamma_\alpha,\gamma_\beta\}=\delta_{\alpha,\beta}\mathrm{I}$ (I the identity operator) and constitute zero energy eigen-operators of the Bogoliubov - de Gennes (BdG) Hamiltonian. Since MFs are hermitian they can be described by the following general expression

$$\gamma_\alpha=\int d\boldsymbol{r}\left[u^*_{\uparrow,\alpha}(\boldsymbol{r})\psi_\uparrow(\boldsymbol{r})+u^*_{\downarrow,\alpha}(\boldsymbol{r})\psi_\downarrow(\boldsymbol{r})+u_{\uparrow,\alpha}(\boldsymbol{r})\psi^\dagger_\uparrow(\boldsymbol{r})+u_{\downarrow,\alpha}(\boldsymbol{r})\psi^\dagger_\downarrow(\boldsymbol{r})\right]\,, \tag{1}$$

where $\psi^\dagger_\sigma(\boldsymbol{r})/\psi_\sigma(\boldsymbol{r})$ correspond to the creation/annihilation operators of an electron in 2d coordinate space $\boldsymbol{r}=(x,y)$ with spin projection $\sigma=\uparrow,\downarrow$. Notice that MFs require linear combinations of electronic operators and their hermitian conjugates. This limits us to systems in which the spin-quantization axis is fixed. Notice that for a system with spin-rotational symmetry, the application of a homogeneous magnetic field breaks the latter symmetry but the spin-quantization axis can be always redefined. In this case, MFs are not accessible. As a matter of fact, MFs can appear only in systems with spatially inhomogeneous magnetism, spin-orbit coupling or spin-triplet superconductivity.

In this work I focus on systems that satisfy the above requirements and are either microscopically or phenomenologically described by the following *bulk* Hamiltonian

$$\begin{aligned}\mathcal{H} &= \int d\boldsymbol{r}\;\hat{\psi}^\dagger(\boldsymbol{r})\left[\frac{\hat{\boldsymbol{p}}^2}{2m}-\mu+V(\boldsymbol{r})-\boldsymbol{M}(\boldsymbol{r})\cdot\boldsymbol{\sigma}+\frac{\{v(\boldsymbol{r}),\hat{p}_x\sigma_y-\hat{p}_y\sigma_x\}}{2}\right]\hat{\psi}(\boldsymbol{r})\\ &+ \int d\boldsymbol{r}\left[\psi^\dagger_\uparrow(\boldsymbol{r})\Delta(\boldsymbol{r})\psi^\dagger_\downarrow(\boldsymbol{r})+\psi_\downarrow(\boldsymbol{r})\Delta^*(\boldsymbol{r})\psi_\uparrow(\boldsymbol{r})\right]\,,\end{aligned} \tag{2}$$

where $\hat{\psi}^\dagger(\boldsymbol{r})=(\psi^\dagger_\uparrow(\boldsymbol{r})\;\;\psi^\dagger_\downarrow(\boldsymbol{r}))$, $\boldsymbol{\sigma}$ are the spin Pauli matrices, $v(\boldsymbol{r})$ is the spatially dependent strength of the Rashba spin-orbit coupling, $\boldsymbol{M}(\boldsymbol{r})$ corresponds to a magnetic-field or a magnetization profile and $\Delta(\boldsymbol{r})$ defines a spatially varying superconducting order parameter. Notice that in some sense the above Hamiltonian is overcomplete, since it covers all the cases that we will consider, without implying that all the terms are simultaneously required for obtaining a TSC. Furthermore, at the level of my topological classification, the origin of the involved terms is unimportant. However, I have to remark that when I will discuss specific cases I will concentrate on engineered TSC, which for instance involve conventional types of magnetism and mainly proximity induced superconductivity [25]. This means that I will not consider here the cases of unconventional [26] density waves [27–31] or superconductors [32], although some of our conclusions could also apply to these systems.

Since for the situations considered in the present study the spin-quantization is always fixed, I will employ the following spinor

$$\widehat{\Psi}^\dagger(\boldsymbol{r})=\Big(\psi^\dagger_\uparrow(\boldsymbol{r})\,,\psi^\dagger_\downarrow(\boldsymbol{r})\,,\psi_\uparrow(\boldsymbol{r})\,,\psi_\downarrow(\boldsymbol{r})\Big)\,, \tag{3}$$

and use the $\boldsymbol{\tau}$ Pauli matrices in order to represent matrices in the Nambu particle-hole space. With the introduction of the above enlarged spinor the Hamiltonian can be rewritten in the following compact way

$$\mathcal{H}=\frac{1}{2}\int d\boldsymbol{r}\;\widehat{\Psi}^{\dagger}(\boldsymbol{r})\widehat{\mathcal{H}}(\hat{\boldsymbol{p}},\boldsymbol{r})\widehat{\Psi}(\boldsymbol{r})\,, \tag{4}$$

where $\widehat{\mathcal{H}}(\hat{\boldsymbol{p}},\boldsymbol{r})$ corresponds to the BdG Hamiltonian. Notice that the factor of 1/2 is crucial for avoiding double counting of the degrees of freedom, since the above spinor does not obey to the usual fermionic commutation relations.

II. TOPOLOGICAL CLASSIFICATION PRINCIPLES

Before discussing the possible topological phases arising from our model Hamiltonian, I will briefly review the basics of how to classify topological systems. My goal is to first highlight a key point which is crucial for classifying TSCs and then demonstrate how this can provide further topological insight concerning previously studied systems [18, 19]. This key point is that topological classification of systems following the recently developed methods [33, 34], is conducted for irreducible Hamiltonians, for which one cannot find any unitary operator $\mathcal{O}_u$ satisfying $[\widehat{\mathcal{H}}(\hat{\boldsymbol{p}},\boldsymbol{r}),\mathcal{O}_u]=0$. If there is a number of these type of operators, we can block diagonalize the Hamiltonian and topologically classify each sub-block. Of course, this is not the only route to study topological properties, since one can also directly construct topological invariants for reducible Hamiltonians [35]. Nevertheless, studying irreducible Hamiltonians provides a transparent analysis of the topological classes. Here I am interested in "hidden" unitary discrete symmetry operators satisfying the cyclic property $\mathcal{O}_u^n=\mathrm{I}$, with $n\in\mathbb{Z}$. In the simplest case $n=2$, we can block diagonalize the Hamiltonian into two sub-blocks labelled by the eigenvalues ±1 of $\mathcal{O}_u$, leading to a direct sum of the form $\widehat{\mathcal{H}}_+(\hat{\boldsymbol{p}},\boldsymbol{r})\oplus\widehat{\mathcal{H}}_-(\hat{\boldsymbol{p}},\boldsymbol{r})$. Notice that because of the discrete symmetry $\mathcal{O}_u$, both sub-blocks are constrained to belong to the same symmetry class. However, the two sub-systems do not necessarily reside in the same topological class.

The symmetry class and the related accessible topological phases of an irreducible Hamiltonian are defined by the possible presence of three specific types of discrete symmetries. The first two correspond to a generalized time-reversal symmetry effected by the anti-unitary operator Θ and a charge conjugation symmetry effected by an anti-unitary operator Ξ. If Θ is a symmetry of the Hamiltonian, it satisfies $[\widehat{\mathcal{H}}(\hat{\boldsymbol{p}},\boldsymbol{r}),\Theta]=0\Rightarrow\Theta^{-1}\widehat{\mathcal{H}}(\hat{\boldsymbol{p}},\boldsymbol{r})\Theta=+\widehat{\mathcal{H}}(\hat{\boldsymbol{p}},\boldsymbol{r})$ while in the case of charge-conjugation we instead have $\{\widehat{\mathcal{H}}(\hat{\boldsymbol{p}},\boldsymbol{r}),\Xi\}=0\Rightarrow\Xi^{-1}\widehat{\mathcal{H}}(\hat{\boldsymbol{p}},\boldsymbol{r})\Xi=-\widehat{\mathcal{H}}(\hat{\boldsymbol{p}},\boldsymbol{r})$. If Θ and Ξ constitute symmetries of the Hamiltonian at the same time, then the Hamiltonian additionaly satisfies $\{\widehat{\mathcal{H}}(\hat{\boldsymbol{p}},\boldsymbol{r}),\Theta\Xi\}=0$ where the combined $\Theta\Xi$ operator is unitary and is termed chiral symmetry operator Π. The inclusion of Π completes the set of symmetries that are required for determining the symmetry class of an irreducible Hamiltonian. In fact, in order to cover all possible symmetry classes, we have to take into account the case in which a unitary chiral symmetry may exist without the necessary presence of Θ and Ξ symmetries.

In this work, I also provide an example of an anti-unitary hidden symmetry $\mathcal{O}_a$. In this case $\mathcal{O}_a$ constitutes an additional generalized time-reversal symmetry which modifies the initial symmetry class of the system, instead of splitting the latter in a direct sum of identical symmetry classes as for the unitary analog $\mathcal{O}_u$. For instance, if a system is initially in class D, then the emergence of an anti-unitary hidden symmetry $\mathcal{O}_a$ with $\mathcal{O}_a^2=+\mathrm{I}$ will change its symmetry class to BDI.

For the cases under consideration, the BdG Hamiltonian enjoys a charge-conjugation symmetry $\Xi=\tau_x\mathcal{K}$, where $\mathcal{K}$ defines complex conjugation. Since $\Xi^2=+\mathrm{I}$, we obtain only the following three allowed symmetry classes presented in Table I: BDI, D, DIII or their direct sums BDI$\oplus$BDI, D$\oplus$D, DIII$\oplus$DIII in the presence of a hidden symmetry $\mathcal{O}_u$. Notice that the classes BDI and DIII are characterized by a time-reversal symmetry Θ with $\Theta^2=+\mathrm{I}$ and $\Theta^2=-\mathrm{I}$, respectively. In the first case, Θ symmetry implies that the Hamiltonian is real while in the second that there exist a Kramers-type degeneracy leading to doublets of solutions. Below I examine the minimal cases that can lead to a symmetry class supporting MFs. For completeness I will also shortly discuss previously studied models.

TABLE I: Symmetry classes of topological superconductors supporting Majorana fermions. For Θ and Ξ, $\pm$I corresponds to the result of Θ^2 and Ξ^2. For Π, I denotes that the symmetry is present. Conversely, 0 implies that the corresponding symmetry is broken.

Class	Θ	Ξ	Π	1d	2d	3d
BDI	+I	+I	I	$\mathbb{Z}$	0	0
D	0	+I	0	$\mathbb{Z}_2$	$\mathbb{Z}$	0
DIII	−I	+I	I	$\mathbb{Z}_2$	$\mathbb{Z}_2$	$\mathbb{Z}$



FIG. 1: Point group symmetries for quasi-1d and strictly 1d geometries of a topological superconductor (depicted with blue). In the quasi-1d case, inversion symmetry along the $z-$axis, σ_h, is broken.

III. RESULTS: POSSIBLE PHASES OF ENGINEERED TOPOLOGICAL SUPERCONDUCTORS

In the present section I carry out a thorough analysis of the accessible TSC phases that follow from the Hamiltonian of Eq. (2). For the strict 2d and 1d cases I will consider that $V(\boldsymbol{r})=0$. To analyze the quasi-1d case, I will always assume the presence of a confining potential $V(y)$. For topological computation applications based on edge MFs the quasi-1d and pure 1d setups are mostly relevant. The possible unitary symmetries that can appear for these systems originate from the point group $\mathcal{G}$ and translation operations $t_{\boldsymbol{a}}$ with $t_{\boldsymbol{a}}\boldsymbol{r}=\boldsymbol{r}+\boldsymbol{a}$. Let me now focus on the point group symmetries for the quasi-1d and pure 1d geometries, which I also graphically demonstrate in Fig. 1. The point group for a quasi-1d system confined in the xy-plane is C_{2v}. This symmetry group includes a C_2 π-rotation about the z-axis ($\boldsymbol{r}\rightarrow-\boldsymbol{r}$, $z\rightarrow z$) and two σ_v reflection operations σ_{xz} ($y\rightarrow-y$) and σ_{yz} ($x\rightarrow-x$), where the indices correspond to the mirroring plane. Notice that the reflection symmetry operation $\sigma_h\equiv\sigma_{xy}$ ($z\rightarrow-z$) is broken in C_{2v}. In the strict 1d case we are left only with σ_{yz}. For random $v(\boldsymbol{r})$, $\boldsymbol{M}(\boldsymbol{r})$ and $\Delta(\boldsymbol{r})$ all the aforementioned symmetries are broken. Nevertheless, for special spatial profiles of the latter functions, a hidden symmetry can emerge, which consists of these basic symmetry operations or other *already broken* symmetries such as σ_h.

In Table II, I present the topological classification for the Hamiltonian of Eq. (2) where all possible unitary symmetries are broken due to the spatial dependence of $v(\boldsymbol{r})$, $\boldsymbol{M}(\boldsymbol{r})$ and $\Delta(\boldsymbol{r})$. I demand that $|\boldsymbol{M}(\boldsymbol{r})|\neq0$ and $|\Delta(\boldsymbol{r})|\neq0$ so as to avoid any gap closings that could lead to a macroscopic coexistence of different topological phases throughout the volume of the material. I also have to remark that in the case of a translationally invariant system, we can transfer to $\boldsymbol{k}-$space in order to calculate topological invariants. If translational symmetry is broken, then analysis of the topological properties in coordinate or momentum space exhibits the same complexity. Of course, there can be also cases where topological properties in combined $(\boldsymbol{r},\boldsymbol{k})$-space can be relevant [35, 36].

One of the most important results is Case II in Table II, where the simultaneous presence of Rashba spin-orbit

TABLE II: Accessible topological superconducting phases due to the combined presence of inhomogeneous Rashba spin-orbit coupling $v(\boldsymbol{r})$, magnetization $\boldsymbol{M}(\boldsymbol{r})$ and superconducting order parameter $\Delta(\boldsymbol{r})$. The presence (absence) of the aforementioned terms is indicated with + (-). The resulting symmetry class depends on the behaviour of the magnetic and superconducting Hamiltonian terms under complex conjugation $\mathcal{K}$, since the Rashba spin-orbit coupling term always preserves $\mathcal{T}$. For a term that is already present (+), we denote the case of preserved (broken) complex conjugation as $\mathcal{K}=\mathrm{I}$ (0). Notice that symmetry classes that lack a strong topological invariant for the corresponding dimensionality are shown with italics and correspond to weak topological superconductors. In the presence of a unitary hidden symmetry $\mathcal{O}_u$ with the cyclic property $\mathcal{O}_u^n=\mathrm{I}$, a symmetry class TC splits into n identical sub-classes $\oplus_n$TC. However, the sub-systems do not have to reside in the same topological class.

Case	$v(\boldsymbol{r})$	$\boldsymbol{M}(\boldsymbol{r})$	$\Delta(\boldsymbol{r})$	2d	quasi-1d	1d
I	$+$	$-$	$\mathcal{K}=\mathrm{I}$	**DIII**	**DIII**	*no MFs*
II	$+$	$-$	$\mathcal{K}=0$	**D**	**D**	*no MFs*
III	$-$	$\mathcal{K}=\mathrm{I}$	$\mathcal{K}=\mathrm{I}$	*BDI*	**BDI**	**BDI**
IV	$-$	$\mathcal{K}=\{0,\mathrm{I},0\}$	$\mathcal{K}=\{\mathrm{I},0,0\}$	**D**	**D**	**D**
V	$+$	$\mathcal{K}=\mathrm{I}$	$\mathcal{K}=\mathrm{I}$	**D**	**D**	**BDI**
VI	$+$	$\mathcal{K}=\{0,\mathrm{I},0\}$	$\mathcal{K}=\{\mathrm{I},0,0\}$	**D**	**D**	**D**

coupling and inhomogeneous superconductivity can lead to MFs in a quasi-1d system, without the requirement of a magnetic field. In fact, an $\boldsymbol{r}$-dependent superconducting phase originating from a supercurrent falls into this case, constituting an experimentally prominent route towards MFs. As far as the table is concerned, the possible phases are essentially classified by the behaviour of the magnetic and superconducting Hamiltonian terms under $\mathcal{K}$.

IV. ANALYSIS OF THE POSSIBLE TOPOLOGICAL PHASES IN THE ABSENCE OF UNITARY SYMMETRIES

In this section I provide the detailed topological classification for the cases presented in the Table II. Notice that for the present discussion the spatial dependence of the terms involved is considered random, unless explicitely stated.

- **Cases** I **and** II

  In the following paragraph I will focus on the Cases I and II that are characterized by the presence of inhomogeneous Rashba spin-orbit coupling $v(\boldsymbol{r})$ and superconducting order parameter $\Delta(\boldsymbol{r})$. The TSCs belonging to these cases are described by the following Hamiltonian

$$\widehat{\mathcal{H}}(\hat{\boldsymbol{p}},\boldsymbol{r})=\left[\frac{\hat{\boldsymbol{p}}^2}{2m}-\mu+V(\boldsymbol{r})\right]\tau_z+\frac{\{v(\boldsymbol{r}),\hat{p}_x\tau_z\sigma_y-\hat{p}_y\sigma_x\}}{2}-\Delta_{\Re}(\boldsymbol{r})\tau_y\sigma_y-\Delta_{\Im}(\boldsymbol{r})\tau_x\sigma_y\,. \tag{5}$$

  The Rashba spin-orbit coupling term is odd under inversion symmetry along the z-axis σ_h, while it is even under the usual time-reversal symmetry $\mathcal{T}$. If the superconducting term is also invariant under $\mathcal{T}$ or equivalently $\mathcal{K}$, since we are dealing with a scalar superconducting order parameter, then $\Delta(\boldsymbol{r})=\Delta_{\Re}(\boldsymbol{r})$ and the full Hamiltonian is characterized by the generalized time-reversal symmetry $\Theta=i\sigma_y\mathcal{K}$ that coincides with $\mathcal{T}$.

  2d system: In the 2d case, the particular system belongs to the symmetry class DIII. Since Θ satisfies $\Theta^2=-\mathrm{I}$, with I the identity operator, we expect boundary MF Kramers doublets. Class DIII possesses a strong $\mathbb{Z}_2$ topological invariant in 2d. The presence of Θ also leads to a chiral symmetry with $\Pi=\tau_x\sigma_y$. In the case where the superconducting order parameter has an additional imaginary component, $\mathcal{T}$ is broken and the system transits to class D. Class D has a strong $\mathbb{Z}$ invariant in 2d and consequently this system constitutes a strong TSC in both cases.

  In order to analyze the symmetry properties in a more transparent manner, I will consider without any loss of generality, the following form for the superconducting order parameter $\Delta(\boldsymbol{r})=\Delta e^{i\boldsymbol{J}\cdot\boldsymbol{r}}$. The particular profile, constitutes the simplest representative of $\mathcal{T}$ violating superconductivity and can be viewed either as the result of the spontaneous formation of a Fulde-Ferrell [37] phase with modulation wave-vector $\boldsymbol{J}$ or the consequence of the application of a supercurrent $\boldsymbol{J}$. The Fulde-Ferrell phase is a special case of pair density waves [38] that have been also recently considered [16] as potential TSCs leading to MFs. On the other hand, the application of supercurrents was previously discussed in Refs. [39]. In the latter implementations a supercurrent was viewed as an extra knob for tuning the topological phase diagram.

  At this point we proceed with gauging away the superconducting phase $\varphi(\boldsymbol{r})=\boldsymbol{J}\cdot\boldsymbol{r}$ via the minimal coupling $\hat{\boldsymbol{p}}\rightarrow\hat{\boldsymbol{p}}+\hbar\boldsymbol{\nabla}\varphi(\boldsymbol{r})\tau_z/2=\hat{\boldsymbol{p}}+\hbar\boldsymbol{J}\tau_z/2$, leading to

$$\begin{aligned}\widehat{\mathcal{H}}'(\hat{\boldsymbol{p}})\;&=\;\frac{\hbar}{2m}\,\boldsymbol{J}\cdot\hat{\boldsymbol{p}}\,I+\left[\frac{\hat{\boldsymbol{p}}^2}{2m}+\frac{(\hbar\boldsymbol{J}/2)^2}{2m}-\mu+V(\boldsymbol{r})\right]\tau_z+\frac{\{v(\boldsymbol{r}),\hat{p}_x\tau_z\sigma_y-\hat{p}_y\sigma_x\}}{2}+\frac{v(\boldsymbol{r})\hbar}{2}\left(J_x\sigma_y-J_y\tau_z\sigma_x\right)\\&-\;\Delta\tau_y\sigma_y\,.\end{aligned} \tag{6}$$

  It is straightforward to confirm that for $\boldsymbol{J}=\boldsymbol{0}$ the system belongs to class DIII because $\mathcal{T}$ is preserved while for finite $\boldsymbol{J}$ the system lies in class D.

  quasi-1d system: In order to investigate the quasi-1d and 1d cases I set $v(\boldsymbol{r})=v(x)$. Furthermore for the quasi-1d case I additionaly switch on a confining potential $V(\boldsymbol{r})=V(y)$. The presence of the confining potential lowers the symmetry of the system, permitting anisotropic coefficients for the Rashba terms $\hat{p}_x\tau_z\sigma_y$ and $\hat{p}_y\sigma_x$, instead of a common $v(x)$. For my analysis I will keep the coefficients equal since the only crucial requirement for my study is that they are both non-zero. To achieve confinement, I consider the case of a harmonic potential $V(y)=m\omega^2y^2/2$. This term is translationally invariant along the x-direction and even under C$_2$, σ_{xz} and σ_{yz}. Another option for the confining potential is the infinite wall potential $V(|y|>L_y)=+\infty$. For the choice of the harmonic confining potential, the Hamiltonian reads

$$\begin{aligned}\widehat{\mathcal{H}}(\hat{p}_x,\hat{a},\hat{a}^\dagger) &= \frac{\hbar}{2m}\left(J_x\hat{p}_x+J_y\sqrt{\frac{m\omega\hbar}{2}}\,\frac{\hat{a}-\hat{a}^\dagger}{i}\right)I+\left[\frac{\hat{p}_x^2}{2m}+\frac{(\hbar\boldsymbol{J}/2)^2}{2m}-\mu+\hbar\omega\left(\hat{a}^\dagger\hat{a}+\frac{1}{2}\right)\right]\tau_z\\ &+ \frac{\{v(x),\hat{p}_x\}}{2}\,\tau_z\sigma_y-v(x)\sqrt{\frac{m\omega\hbar}{2}}\,\frac{\hat{a}-\hat{a}^\dagger}{i}\sigma_x+\frac{v(x)\hbar}{2}\left(J_x\sigma_y-J_y\tau_z\sigma_x\right)-\Delta\tau_y\sigma_y\,,\end{aligned}\tag{7}$$

where I introduced the quantum harmonic oscillator's bosonic creation (annihilation) operator $\hat{a}^\dagger$ ($\hat{a}$). By introducing the eigenfunctions $|n\rangle$ of the number operator $\widehat{N}=\hat{a}^\dagger\hat{a}$, I obtain the matrix Hamiltonian

$$\begin{aligned}\widehat{\mathcal{H}}(\hat{p}_x) &= \frac{\hbar}{2m}\left(J_x\hat{p}_xI+J_y\hbar\lambda_y\right)+\left(\frac{\hat{p}_x^2}{2m}-\mu\lambda_z^{\boldsymbol{J}}\right)\tau_z+\frac{\{v(x),\hat{p}_x\}}{2}\,\tau_z\sigma_y-v(x)\hbar\lambda_y\sigma_x+\frac{v(x)\hbar}{2}\left(J_x\sigma_y-J_y\tau_z\sigma_x\right)\\ &- \Delta\tau_y\sigma_y\,,\end{aligned}\tag{8}$$

that is defined in spin, Nambu and $\widehat{N}$ spaces with

$$\langle n|\,\lambda_z^{\boldsymbol{J}}\,|s\rangle=\delta_{n,s}\left[\mu-\frac{(\hbar\boldsymbol{J}/2)^2}{2m}-\hbar\omega\left(n+\frac{1}{2}\right)\right]/\mu\quad\text{and}\quad\langle n|\,\lambda_y\,|s\rangle=\sqrt{\frac{m\omega}{2\hbar}}\,\frac{\sqrt{n+1}\delta_{n,s-1}-\sqrt{n}\delta_{n,s+1}}{i}\,.\tag{9}$$

Since the form of the Hamiltonian is identical to the 2d case and $\mathcal{K}^{-1}\lambda_y\mathcal{K}=-\lambda_y$ (similarly to $\hat{p}_y$), the quasi-1d model also belongs to the DIII class for $\boldsymbol{J}=\boldsymbol{0}$ and to class D for $\boldsymbol{J}\neq\boldsymbol{0}$.

1d system: For studying the strictly 1d system, I apply the dimensional reduction method to the 2d model of Eq. (6) and set $\hat{p}_y=J_y=0$, that yields

$$\widehat{\mathcal{H}}'(\hat{p}_x)=\frac{\hbar}{2m}\,J_x\hat{p}_xI+\left[\frac{\hat{p}_x^2}{2m}+\frac{\left(\hbar J_x/2\right)^2}{2m}-\mu\right]\tau_z+\frac{\{v(x),\hat{p}_x\}}{2}\,\tau_z\sigma_y+\frac{v(x)}{2}\,\hbar J_x\sigma_y-\Delta\tau_y\sigma_y\,.\tag{10}$$

We observe that for this model we retain our freedom to redefine the spin-quantization axis and as a result the above Hamiltonian does not support MFs.

- **Cases** III **and** IV

In this section I consider TSC phases that do not involve spin-orbit coupling. This implies that at least two components of an inhomogeneous magnetization field must be present in order to lock the spin-quantization axis, since the latter constitutes a prerequisite for obtaining MFs. For this kind of systems, the Hamiltonian reads

$$\widehat{\mathcal{H}}(\hat{\boldsymbol{p}},\boldsymbol{r})=\left[\frac{\hat{\boldsymbol{p}}^2}{2m}-\mu+V(\boldsymbol{r})\right]\tau_z-\boldsymbol{M}(\boldsymbol{r})\cdot(\tau_z\sigma_x,\sigma_y,\tau_z\sigma_z)-\Delta_{\Re}(\boldsymbol{r})\tau_y\sigma_y-\Delta_{\Im}(\boldsymbol{r})\tau_x\sigma_y\,.\tag{11}$$

For the specific type of TSCs, the magnetization field $\boldsymbol{M}(\boldsymbol{r})$ is odd under the usual time-reversal symmetry $\mathcal{T}$. However, its behavior under complex conjugation $\mathcal{K}$ is not fixed. If $M_y(\boldsymbol{r})=0$ then $\boldsymbol{M}(\boldsymbol{r})$ preserves $\mathcal{K}$. This leads to the following two possiblities depending also on the behaviour of the superconducting order parameter under $\mathcal{K}$. In the first possibility the magnetic and superconducting terms are simultaneously invariant under $\mathcal{K}$ and a generalized time-reversal symmetry appears with $\Theta=\mathcal{K}$ accompanied by a chiral symmetry $\Pi=\tau_x$.

2d system: In 2d, the system belongs to the BDI class that however is not characterized by a strong topological invariant for this dimensionality. Consequently, the specific system corresponds to a weak TSC, since under special circumstances one can generally define weak invariants. The second possibility involves the breaking of $\mathcal{K}$ by either one of the terms. In the latter case, the Hamiltonian belongs to class D which has a strong $\mathbb{Z}$ topological invariant in 2d.

quasi-1d system: For the particular study I will consider for convenience that $\Delta(\boldsymbol{r})=\Delta e^{i\boldsymbol{J}\cdot\boldsymbol{r}}$. As previously, I gauge away the superconducting phase and obtain the equivalent model

$$\widehat{\mathcal{H}}'(\hat{\boldsymbol{p}})=\frac{\hbar}{2m}\,\boldsymbol{J}\cdot\hat{\boldsymbol{p}}\,I+\left[\frac{\hat{\boldsymbol{p}}^2}{2m}+\frac{\left(\hbar\boldsymbol{J}/2\right)^2}{2m}-\mu+V(\boldsymbol{r})\right]\tau_z-\boldsymbol{M}(\boldsymbol{r})\cdot(\tau_z\sigma_x,\sigma_y,\tau_z\sigma_z)-\Delta\tau_y\sigma_y\,.\tag{12}$$

For effecting confinement I will employ once again a harmonic oscillator's potential $V(y) = m\omega^2 y^2/2$ and we also have $\boldsymbol{M}(\boldsymbol{r}) = \boldsymbol{M}(x, \hat{a} + \hat{a}^\dagger)$. Following the same steps as in Cases I and II, I obtain the Hamiltonian

$$\widehat{\mathcal{H}}'(\hat{p}_x) = \frac{\hbar}{2m}\left(J_x \hat{p}_x I + J_y \hbar \lambda_y\right) + \left(\frac{\hat{p}_x^2}{2m} - \mu \lambda_z^{\boldsymbol{J}}\right)\tau_z - \widehat{\boldsymbol{M}}(x)\cdot(\tau_z\sigma_x, \sigma_y, \tau_z\sigma_z) - \Delta\tau_y\sigma_y\,, \tag{13}$$

where $\widehat{\boldsymbol{M}}(x)$ is a symmetric matrix defined in $|n\rangle$ space. If $\boldsymbol{J} = \boldsymbol{0}$ and $\widehat{M}_y(x) = 0$, $\mathcal{K}$ is a symmetry of the Hamiltonian and the system belongs to class BDI. Instead, if $\widehat{M}_y(x) \neq 0$, the system belongs to class D. For the special case where $\boldsymbol{M}(\boldsymbol{r})$ does not depend on the y-coordinate, i.e. $\boldsymbol{M}(\boldsymbol{r}) = \boldsymbol{M}(x)$, $\boldsymbol{M}(\boldsymbol{r})$ becomes diagonal and can be divided into an infinite number of sub-spaces labelled by n yielding

$$\widehat{\mathcal{H}}_n'(\hat{p}_x) = \left[\frac{\hat{p}_x^2}{2m} - \mu + \hbar\omega\left(n + \frac{1}{2}\right)\right]\tau_z - \boldsymbol{M}(x)\cdot(\tau_z\sigma_x, \sigma_y, \tau_z\sigma_z) - \Delta\tau_y\sigma_y\,, \tag{14}$$

which leads to the total symmetry class $\oplus_n$BDI. If $\boldsymbol{M}(x)$ violates $\mathcal{K}$ we obtain a direct sum $\oplus_n$D. By allowing a finite $\boldsymbol{J}$ we also violate $\mathcal{K}$. Specifically, if $\boldsymbol{J} = (J_x, 0)$, that corresponds to the case $\Delta(\boldsymbol{r}) = \Delta(x)$, the system resides in the class $\oplus_n$D. However, if $\boldsymbol{J} = (0, J_y)$ the system belongs to class D, due to the simultaneous presence of λ_y and $\lambda_z^{\boldsymbol{J}}$ in the Hamiltonian of Eq. (13), that does not allow the decomposition in n-sectors. In Table II the general case where $\boldsymbol{M}$ and Δ depend on both coordinates is depicted.

1d system: By dimensional reduction on the Hamiltonian of Eq. (12) we obtain the following pure 1d model

$$\widehat{\mathcal{H}}'(\hat{p}_x) = \frac{\hbar}{2m}\,J_x\hat{p}_x I + \left[\frac{\hat{p}_x^2}{2m} + \frac{\left(\hbar J_x/2\right)^2}{2m} - \mu\right]\tau_z - \boldsymbol{M}(x)\cdot(\tau_z\sigma_x, \sigma_y, \tau_z\sigma_z) - \Delta\tau_y\sigma_y\,. \tag{15}$$

If $M_y(x) = 0$ and $J_x = 0$, $\mathcal{K}$ is conserved and the system belongs to class BDI. Instead, if one of the previous terms is non-zero, the Hamiltonian is not real any more and it falls into symmetry class D [17].

- **Cases** V **and** VI

In the last part of this section I complete the possible cases by considering the situation where all the terms of Eq. (2) are present. The latter equation in combined Nambu and spin spaces reads

$$\begin{aligned}\widehat{\mathcal{H}}(\hat{\boldsymbol{p}}, \boldsymbol{r}) \;=\;& \left[\frac{\hat{\boldsymbol{p}}^2}{2m} - \mu + V(\boldsymbol{r})\right]\tau_z + \frac{\{v(\boldsymbol{r}), \hat{p}_x\tau_z\sigma_y - \hat{p}_y\sigma_x\}}{2} - \boldsymbol{M}(\boldsymbol{r})\cdot(\tau_z\sigma_z, \sigma_y, \tau_z\sigma_z) - \Delta_{\Re}(\boldsymbol{r})\tau_y\sigma_y \\ -\;& \Delta_{\Im}(\boldsymbol{r})\tau_x\sigma_y\,.\end{aligned} \tag{16}$$

When magnetism and Rashba spin-orbit coupling coexist, the accessible topological phases constitute an overlap of the previously examined separate cases. Therefore here we will investigate what are the consequences of the addition of magnetism in Cases I and II for different dimensionalities. Earlier, we observed that when magnetism is not present, there are two possible scenarios depending on the behaviour of the superconducting order parameter under $\mathcal{K}$.

2d and quasi-1d systems: For the specific cases, if $\Delta(\boldsymbol{r}) = \Delta_{\Re}(\boldsymbol{r})$ the system resides in the symmetry class DIII being invariant under $\mathcal{T}$. If $\boldsymbol{M}(\boldsymbol{r})$ is introduced, $\mathcal{T}$ will be broken and the system will transit to class D. If $\Delta(\boldsymbol{r})$ is complex, the system is already in class D, and consequently the inclusion of magnetism leads to no additional effects.

1d system: For pure 1d systems the presence of a magnetic order is crucial and leads to new TSC phases. The 1d descendant of the above Hamiltonian reads

$$\widehat{\mathcal{H}}(\hat{p}_x, x) = \left(\frac{\hat{p}_x^2}{2m} - \mu\right)\tau_z + \frac{\{v(x), \hat{p}_x\}}{2}\,\tau_z\sigma_y - \boldsymbol{M}(x)\cdot(\tau_z\sigma_z, \sigma_y, \tau_z\sigma_z) - \Delta_{\Re}(x)\tau_y\sigma_y - \Delta_{\Im}(x)\tau_x\sigma_y\,. \tag{17}$$

From Table II we immediately observe that there are no TSC phases in the absence of magnetism. As mentioned earlier, the reason is that the presence of the spin-orbit coupling term $\tau_z\sigma_y$ alone, cannot fix the spin-quantization axis. Nevertheless, the addition of a perpendicular magnetization field remedies this problem and can lead to TSC phases. If $\boldsymbol{M}(x)$ is invariant under $\mathcal{K}$, i.e. $M_y(x) = 0$, the Hamiltonian is characterized by a generalized time-reversal symmetry $\Theta = \mathcal{K}$ and a chiral symmetry $\Pi = \tau_x$ which permits an integer number of MFs per edge [41]. The translationally invariant version of this model

$$\widehat{\mathcal{H}}(\hat{p}_x) = \left(\frac{\hat{p}_x^2}{2m} - \mu\right)\tau_z + v\hat{p}_x\tau_z\sigma_y - \boldsymbol{M}\cdot(\tau_z\sigma_z, \sigma_y, \tau_z\sigma_z) - \Delta\tau_y\sigma_y\,, \tag{18}$$

corresponds to the celebrated MF-wire proposal [5] which currently under intense experimental investigation [2] and concerns a Rashba semiconducting wire in the presence of a Zeeman field and proximity induced superconductivity. The systems transits to the topologically non-trivial phase when the criterion

$$|\boldsymbol{M}| > \sqrt{\mu^2 + \Delta^2}\,, \tag{19}$$

is satisfied. Finally, if $M_y(x) \neq 0$ then $\mathcal{K}$ is broken and the system belongs to class D with a $\mathbb{Z}_2$ invariant allowing for a single MF per edge.

V. EXAMPLES OF TOPOLOGICAL PHASES WITH HIDDEN SYMMETRIES

In this section I will demonstrate two examples where unitary or anti-unitary hidden symmetries occur for some of the TSC phases presented in Table II and demonstrate what are the concomitant modifications of the initial symmetry class.

- **Cases** I **and** II **in the presence of a single unitary hidden symmetry** $\boldsymbol{\mathcal{O}}_u$

  Let us now investigate the consequences of the emergence of a "hidden" symmetry due to the special form of the Rashba spin-orbit coupling term. As a case study I will focus on the topological properties of the following quasi-1d Hamiltonian (in 1d we do not obtain a TSC), introduced in Eq. (8) of Case I

$$\begin{aligned}\widehat{\mathcal{H}}(\hat{p}_x) &= \frac{\hbar}{2m}\left(J_x\hat{p}_x I + J_y\hbar\lambda_y\right) + \left(\frac{\hat{p}_x^2}{2m} - \mu\lambda_z^{\boldsymbol{J}}\right)\tau_z + \frac{\{v(x),\hat{p}_x\}}{2}\,\tau_z\sigma_y - v(x)\hbar\lambda_y\sigma_x + \frac{v(x)\hbar}{2}\left(J_x\sigma_y - J_y\tau_z\sigma_x\right)\\ &- \Delta\tau_y\sigma_y\,. \end{aligned}\tag{20}$$

  Here we will restrict to the special situation where $t_{\pi/Q}v(x) = v(x+\pi/Q) = -v(x)$. We may readily observe in which manner this property leads to an emergent unitary symmetry. The terms of the Hamiltonian that do not contain $v(x)$ are invariant under arbitrary translations and under the action of σ_h, which in our formalism is represented as $\sigma_h = i\tau_z\sigma_z$ in spin-space. Since all the terms with coefficient $v(x)$ are odd under σ_h, the full Hamiltonian is invariant under the action of $\mathcal{O}_u = \sigma_h t_{\pi/Q}$. The appearance of a hidden symmetry $\mathcal{O}_u$ leads to an additional generalized time-reversal symmetry $\widetilde{\Theta} = \mathcal{O}_u\mathcal{T}$ and a concomitant chiral symmetry $\widetilde{\Pi} = \mathcal{O}_u\Pi$. The emergence of $\mathcal{O}_u$ modifies the symmetry class of the system by splitting the symmetry classes DIII and D found earlier, into DIII⊕DIII and D⊕D, respectively. Notice that the possibility of point group symmetry protected phases has been recently discussed [42]. A simple example for $v(x+\pi/Q) = -v(x)$ is $v(x) = 2v_Q\cos(Qx+\theta)$ with $Q = 2q$. Here, θ is considered pinned to a constant value. The modulated spin-orbit coupling term can be viewed as an unconventional spin triplet density wave [27], similar to the Rashba spin-orbit density wave proposed in [43] as a potential candidate for the so called "hidden order", which appears in the non-superconducting regime of the phase diagram of the heavy fermion compound URu$_2$Si$_2$.

  The simultaneous presence of the momentum operator $\hat{p}_x$ and coordinate x does not allow for a direct and transparent inspection of further topological properties of the system. Nonetheless, it is also possible in principle to obtain through some kind of "deformation" procedure (in the topological sense) a model defined solely in momentum space that shares the same symmetries and topological properties with the original model. In order for this mapping to be meaningful and offer a direct computation of topological invariants, translational symmetry must be somehow restored. The presence of a periodic $v(x) = 2v_Q\cos(Qx+\theta)$ term, leads to the formation of a band structure with a Brillouin zone of length Q since $t_{2\pi/Q}v(x) = v(x)$. The property $t_{\pi/Q}v(x) = -v(x)$ gives rise to a sublattice structure that will eventually lead to the two sub-blocks of the Hamiltonian that become relevant in the presence of $\mathcal{O}_u$. Since we are not interested in the full band structure, but mainly in restoring translational symmetry, we may expand the field operator in the following fashion

$$\hat{\psi}(x) \simeq e^{+iqx}\hat{\psi}_{+q}(x) + e^{-iqx}\hat{\psi}_{-q}(x)\,, \tag{21}$$

where $\hat{\psi}_{\pm q}(x)$ are slowly varying fields leading to the Hamiltonian

$$\begin{aligned}\widehat{\mathcal{H}}_q(\hat{p}_x) \;=\; & \frac{\hbar}{2m}\left(J_x\hat{p}_x I + J_y\hbar\lambda_y\right) + \frac{\hbar}{2}\tilde{v}J_x\rho_z + \left[\frac{\hat{p}_x^2}{2m} + \frac{(\hbar q)^2}{2m} - \mu\lambda_z^{\boldsymbol{J}}\right]\tau_z + \tilde{v}\hat{p}_x\tau_z\rho_z \\ & + v_Q\hat{p}_x\left(\cos\theta\tau_z\rho_x\sigma_y - \sin\theta\tau_z\rho_y\sigma_y\right) - v_Q\hbar\left(\cos\theta\rho_x\lambda_y\sigma_x - \sin\theta\rho_y\lambda_y\sigma_x\right) \\ & + \frac{v_Q\hbar}{2}J_x\left(\cos\theta\rho_x\sigma_y - \sin\theta\rho_y\sigma_y\right) - \frac{v_Q\hbar}{2}J_y\left(\cos\theta\tau_z\rho_x\sigma_x - \sin\theta\tau_z\rho_y\sigma_x\right) - \Delta\tau_y\sigma_y\,, \end{aligned} \tag{22}$$

with $\tilde{v} = \hbar q/m$. The above Hamiltonian acts on the enlarged spinor

$$\widehat{\Psi}_q^\dagger(x) = \Big(\psi_{+q\uparrow}^\dagger(x)\,,\psi_{+q\downarrow}^\dagger(x)\,,\psi_{-q\uparrow}^\dagger(x)\,,\psi_{-q\downarrow}^\dagger(x)\,,\psi_{-q\uparrow}(x)\,,\psi_{-q\downarrow}(x)\,,\psi_{+q\uparrow}(x)\,,\psi_{+q\downarrow}(x)\Big)\,, \tag{23}$$

with the $\boldsymbol{\rho}$ Pauli matrices acting on $\pm q$ space. Notice that terms with ρ_x or ρ_y carry momentum Q. By expanding the field operator in this manner, we managed to end up with a translationally invariant Hamiltonian. This approximation allows us to readily study topological aspects in momentum space which for the specific case is an easier task compared to the required analysis in coordinate space. Nonetheless, it is not a priori ensured that the coordinate and momentum pictures are topologically equivalent. If they do, this approximation constitutes a suitable deformation procedure for mapping x to k_x space topology.

In order to confirm if these systems belong to the same symmmetry class, we have to study the emerging symmetries for the latter model. In this basis the expression for the generalized time-reversal symmetry operator simplifies to $\Theta = i\sigma_y\mathcal{K} = i\rho_x\sigma_y\mathcal{K}'$, where $\mathcal{K}'$ is a complex conjugation operator not acting on Q or q. The presence of ρ_x effects complex conjugation operation $\mathcal{K}$ in q-space, since $q \to -q$ is equivalent to $\mathcal{K} = \rho_x\mathcal{K}'$. Furthermore, within the specific framework $t_{\pi/Q} = -i\rho_z$, up to an unimportant $\boldsymbol{k}$-dependent phase. Notice that $(-i\rho_z)^2 = -\mathrm{I}$, i.e. similar to the behaviour of rotation operators for a spin-1/2. This is a direct consequence of the fact that the spinor contains the wave-vector q which is half of the wave-vector $Q = 2q$. We may directly confirm that $[\widehat{\mathcal{H}}_q(\hat{p}_x), \sigma_h t_{\pi/Q}] \equiv [\widehat{\mathcal{H}}_q(\hat{p}_x), \tau_z\rho_z\sigma_z] = 0$. When $\boldsymbol{J} = \boldsymbol{0}$, the Hamiltonian is invariant under Θ and the presence of $\mathcal{O}_u = \sigma_h t_{\pi/Q}$ leads to the additional time-reversal symmetry $\widetilde{\Theta} = \mathcal{O}_u\Theta = i\tau_z\rho_y\sigma_x\mathcal{K}'$. The emerging chiral symmetries for this model read $\Pi = \tau_x\sigma_y$ and $\widetilde{\Pi} = \tau_y\rho_z\sigma_x$. As expected, in this case the system belongs to the symmetry class DIII$\oplus$DIII. Furthermore, if $\boldsymbol{J} \neq \boldsymbol{0}$ then Θ is broken and the system transits to the symmetry class D$\oplus$D. To explicitely demonstrate the sub-block structure of the Hamiltonian and the direct sum of symmetry classes, I effect the unitary transformation

$$\Lambda = \frac{\mathrm{I} + \rho_z\sigma_z}{2} + \tau_x\frac{\mathrm{I} - \rho_z\sigma_z}{2}\,, \tag{24}$$

which transforms the Hamiltonian as follows $\widehat{\mathcal{H}}_q^{rot}(\hat{p}_x) = \Lambda^\dagger\widehat{\mathcal{H}}_q(\hat{p}_x)\Lambda$. This particular unitary transformation block diagonalizes the matrix $\tau_z\rho_z\sigma_z$, representing the hidden symmetry operation $\mathcal{O}_u$, into $\Lambda^\dagger\tau_z\rho_z\sigma_z\Lambda = \tau_z$. The transformed Hamiltonian is block diagonal and can be labelled by the eigenvalues of τ_z, $\tau = \pm 1$, yielding

$$\begin{aligned}\widehat{\mathcal{H}}_{q,\tau}^{rot}(\hat{p}_x) \;=\; & \frac{\hbar}{2m}\left(J_x\hat{p}_x + J_y\hbar\lambda_y\right)I + \frac{\hbar}{2}\tilde{v}J_x\rho_z + \tau\left[\frac{\hat{p}_x^2}{2m} + \frac{(\hbar q)^2}{2m} - \mu\lambda_z^{\boldsymbol{J}}\right]\rho_z\sigma_z + \tau\tilde{v}\hat{p}_x\sigma_z \\ & + v_Q\left(\tau\hat{p}_x\cos\theta + \hbar\lambda_y\sin\theta\right)\rho_y\sigma_x + v_Q\left(\tau\hat{p}_x\sin\theta - \hbar\lambda_y\cos\theta\right)\rho_x\sigma_x \\ & + \frac{v_Q\hbar}{2}\left[\left(J_x\cos\theta + \tau J_y\sin\theta\right)\rho_x\sigma_y - \left(J_x\sin\theta - \tau J_y\cos\theta\right)\rho_y\sigma_y\right] + \tau\Delta\rho_z\sigma_x\,. \end{aligned} \tag{25}$$

I have to remark that the above topological classification conclusions hold for a bulk system. In order to observe the two sets of edge MFs one must introduce boundaries that preserve the $\mathcal{O}_u$ symmetry. Notice however that in the present example the preservation of $\mathcal{O}_u$ requires a specific behaviour under the translation operation $t_{\pi/Q}$, which will be unavoidably broken for a finite system. Nonetheless, the case discussed here does not only constitute an example of mere academic interest. Although the presence of protected boundary modes [3, 44] is considered to be the hallmark of topologically non-trivial phases, it does not constitute the unique route for diagnosing topological order. In fact, fingerprints of topological non-trivial phases can be also found in manifestations of the bulk system. One example is the polar Kerr effect [45] that characterizes class D chiral p-wave superconductors. This experiment can provide a direct evidence of topological order by solely probing



FIG. 2: Engineered topological superconductor consisting of a metal with spin-spiral magnetic order $\boldsymbol{M}(x) = \big(2M_x^Q\cos Qx, 0, 2M_z^Q\sin Qx\big)$ placed on top of a bulk conventional superconductor. The magnetic order is invariant under the combined action of time-reversal symmetry $\mathcal{T}$ followed by a π/Q translation operation, $t_{\pi/Q}$. The presence of the anti-unitary hidden symmetry $\mathcal{O}_a = t_{\pi/Q}\mathcal{T}$ leads to the class BDI⊕BDI, compared to class BDI when it is absent.

the bulk response. Similar chiral phenomena emerge in non-superconducting systems. In the latter, apart from a similar polar Kerr effect [29, 46], an anomalous thermoelectric Nernst response [30, 47] and a topological Meissner effect [31] constitute additional smoking gun signatures of quantum anomalous Hall phases (class A). In fact, topological response survives also in finite temperatures, though exhibiting no quantization phenomena. Evenmore, the bulk magnetic response [48] of a quantum spin Hall insulator (class AII) can provide alternative routes for confirming the transition to the topologically non-trivial phase.

- **Cases** III **and** IV **with a single anti-unitary hidden symmetry $\mathcal{O}_a$**

Here I will investigate the consequences of the emergence of an anti-unitary hidden symmetry $\mathcal{O}_a$ on the symmetry class of the 1d model of Eq. (15) that was obtained for Cases III and IV. In this way I will be in a position to make a connection to previous studies [18, 19]. The Hamiltonian of Eq. (15) for $J_x = 0$, has the following form

$$\widehat{\mathcal{H}}(\hat{p}_x, x) = \left(\frac{\hat{p}_x^2}{2m} - \mu\right)\tau_z - \boldsymbol{M}(x)\cdot(\tau_z\sigma_x, \sigma_y, \tau_z\sigma_z) - \Delta\tau_y\sigma_y\,. \tag{26}$$

First I will focus on the Case III, which belongs to class BDI with $\Theta = \mathcal{K}$, if $\boldsymbol{M}(x)$ is random and $M_y(x) = 0$. I demonstrate that the topological properties of the system change with the emergence of a hidden symmetry. I now assume that $t_{\pi/Q}\boldsymbol{M}(x) = \boldsymbol{M}(x+\pi/Q) = -\boldsymbol{M}(x)$. For this special case, the Hamiltonian is invariant under the anti-unitary hidden symmetry $\mathcal{O}_a = t_{\pi/Q}\mathcal{T}$. Since $t_{\pi/Q}$ is unitary and $\mathcal{T}$ anti-unitary, the particular symmetry constitutes an additional generalized time-reversal symmetry $\widetilde{\Theta} = \mathcal{O}_a$. In order to gain more insight, I will consider the simple spin-spiral magnetization profile $M_x(x) = 2M_x^Q\cos Qx$ and $M_z(x) = 2M_z^Q\sin Qx$ which is depicted in Fig. 2. Prior studies [18, 19] have focused on the case $M_x^Q = M_z^Q$. By expanding the field operator as in Eq. (21) we obtain

$$\widehat{\mathcal{H}}(\hat{p}_x) = \left[\frac{\hat{p}_x^2}{2m} + \frac{(\hbar q)^2}{2m} - \mu\right]\tau_z + \tilde{\upsilon}\hat{p}_x\tau_z\rho_z - M_x^Q\tau_z\rho_x\sigma_x + M_z^Q\tau_z\rho_y\sigma_z - \Delta\tau_y\sigma_y\,. \tag{27}$$

Within this framework we have $\Theta = \mathcal{K} = \rho_x\mathcal{K}'$ and $\widetilde{\Theta} = \mathcal{O}_a = t_{\pi/Q}\mathcal{T} = -i\rho_z i\sigma_y\mathcal{K} = -i\rho_z i\sigma_y\rho_x\mathcal{K}' = i\rho_y\sigma_y\mathcal{K}'$. Remember that $\mathcal{K}'$ does not act on q and Q. We readily observe that for both generalized time-reversal symmetries we have $\Theta^2 = \widetilde{\Theta}^2 = +\mathrm{I}$ leading to the symmetry class BDI⊕BDI. Note that in Refs. [18, 19] it was shown that, up to a spatially dependent unitary transformation, the model of Eq. (26) is equivalent to the MF-wire model of Eq. (18) with the latter belonging to the symmetry class BDI. However, as we showed here the particular system belongs to class BDI⊕BDI. Generally a unitary equivalence does not necessarily constitute a topological equivalence. Nonetheless, it is straighforward to demonstrate that the two pictures agree with each other. By performing the transformation $\widehat{\mathcal{H}}^{rot}(\hat{p}_x) = \Lambda^\dagger\widehat{\mathcal{H}}^{rot}(\hat{p}_x)\Lambda$ with $\Lambda = (\rho_z\sigma_z + \sigma_y)(\rho_y + \rho_z)/2$ we obtain

$$\widehat{\mathcal{H}}^{rot}(\hat{p}_x) = \left[\frac{\hat{p}_x^2}{2m} + \frac{(\hbar q)^2}{2m} - \mu\right]\tau_z + \tilde{\upsilon}\hat{p}_x\tau_z\rho_y + M_x^Q\tau_z\rho_z - M_z^Q\tau_z\rho_z\sigma_z - \Delta\tau_y\rho_y\sigma_z\,. \tag{28}$$

As anticipated, the above Hamiltonian is block diagonal and if we introduce the eigenstates $\sigma = \pm 1$ of σ_z in the rotated frame, we obtain

$$\widehat{\mathcal{H}}_{\sigma}^{rot}(\hat{p}_x) = \left[\frac{\hat{p}_x^2}{2m} + \frac{(\hbar q)^2}{2m} - \mu\right]\tau_z + \tilde{v}\hat{p}_x\tau_z\rho_y + \left(M_x^Q - \sigma M_z^Q\right)\tau_z\rho_z - \sigma\Delta\tau_y\rho_y\,. \tag{29}$$

Remarkably, each of the above block Hamiltonians is identical to the MF-wire Hamiltonian of Eq. (18) with effective chemical potential $\mu - (\hbar q)^2/2m$, spin-orbit coupling strength $\tilde{v}$, Zeeman field $M_x^Q - \sigma M_z^Q$ and superconducting order parameter $\sigma\Delta$. Each of the blocks will be in the topologically non-trivial phase when the criterion

$$\left|M_x^Q - \sigma M_z^Q\right| > \sqrt{\left[\mu - (\hbar q)^2/2m\right]^2 + \Delta^2}\,, \tag{30}$$

is satisfied in complete analogy to Eq. (19). We observe that the two sub-systems are not necessarily in the topologically non-trivial phase, at the same time. Evenmore, if we consider $M_x^Q = M_z^Q$ as in prior studies [18, 19], only one of the sub-systems can be in the topologically non-trivial phase. In this case, the system effectively behaves as a class BDI TSC and this is in accordance with the previous analytical findings. Notice that for a bounded system the hidden symmetry will be broken and the multiple edge MFs are not expected to be observed even when $M_x^Q \neq M_z^Q$.

VI. NEW TQC PERSPECTIVES IN A TSC WITH UNITARY HIDDEN DISCRETE SYMMETRIES

For the cases that we considered in this work, hidden symmetry involved a specific behaviour under translations. However, this type of symmetry will be explicitely broken when boundaries are introduced. Nevertheless, one could look for alternative, robust and tunable, hidden symmetries that are related to some internal degree of freedom such as a valley, orbital or band index.

Let us now discuss new routes that open up for topological quantum computing when we consider the additional presence of a hidden unitary discrete symmetry $\mathcal{O}_u$. Generally, two edge MFs γ_a and γ_b combine into a zero-energy fermionic mode $d = (\gamma_a + i\gamma_b)/\sqrt{2}$ that leads to a doubly degenerate ground state $|1\rangle$ and $|0\rangle$. These two states correspond to many-body ground states where the zero-mode is occupied or not, respectively. The Hilbert space spanned by these two degenerate states defines a topological qubit which is in principle [49] protected by decoherence due to the non-local binding of the MFs. The fundamental and in fact the only allowed topologically protected single qubit operation which we may perform is called braiding [6, 7] and corresponds to exchanging the two MFs in real space Fig. 3. After braiding is effected the states $|1\rangle$ and $|0\rangle$ become $e^{i\pi/4}|1\rangle$ and $e^{-i\pi/4}|0\rangle$, picking up a relative $\pi/2$ phase. For a counterclockwise rotation, braiding corresponds to the transformation $\gamma_a \to +\gamma_b$ and $\gamma_b \to -\gamma_a$ while for a clockwise rotation we obtain the inverse transformation $\gamma_a \to -\gamma_b$ and $\gamma_b \to +\gamma_a$ [7].



FIG. 3: Exchanging Majorana fermions in real space leads to a protected single topological qubit rotation termed braiding. In qubit space, this corresponds to a gate operation where the qubit states pick up a relative $\pi/2$ phase.

For a system that supports a single MF per edge, the emergence of a unitary discrete symmetry $\mathcal{O}_u$ with $\mathcal{O}_u^2 = \mathrm{I}$, can lead to an additional MF per edge. The two MFs per edge are labelled by the ± 1 eigenstates of $\mathcal{O}_u$, leading to the following 4 edge MFs $\gamma_{a\pm}$ and $\gamma_{b\pm}$. With the latter, we can define two zero-modes $d_\pm = (\gamma_{a\pm} + i\gamma_{b\pm})/\sqrt{2}$ and two topological qubits with states $\{|1_+\rangle, |0_+\rangle\}$ and $\{|1_-\rangle, |0_-\rangle\}$. The accessible protected non-Abelian operations that we may perform within this four-fold degenerate Hilbert space are restricted by the *simultaneous conservation of fermion parity and* $\mathcal{O}_u$. Essentially, the possible operations are combinations of simultaneous or separate clockwise and counterclockwise braiding operations in each of the topological qubit spaces. Specifically, we have the following four operations:

$$A:\{|1_+\rangle,|0_+\rangle,|1_-\rangle,|0_-\rangle\} \;\rightarrow\; \{e^{i\pi/4}|1_+\rangle,e^{-i\pi/4}|0_+\rangle,|1_-\rangle,|0_-\rangle\}\,, \tag{31}$$

$$B:\{|1_+\rangle,|0_+\rangle,|1_-\rangle,|0_-\rangle\} \;\rightarrow\; \{|1_+\rangle,|0_+\rangle,e^{i\pi/4}|1_-\rangle,e^{-i\pi/4}|0_-\rangle\}\,, \tag{32}$$

$$C:\{|1_+\rangle,|0_+\rangle,|1_-\rangle,|0_-\rangle\} \;\rightarrow\; \{e^{i\pi/4}|1_+\rangle,e^{-i\pi/4}|0_+\rangle,e^{i\pi/4}|1_-\rangle,e^{-i\pi/4}|0_-\rangle\}\,, \tag{33}$$

$$D:\{|1_+\rangle,|0_+\rangle,|1_-\rangle,|0_-\rangle\} \;\rightarrow\; \{e^{i\pi/4}|1_+\rangle,e^{-i\pi/4}|0_+\rangle,e^{-i\pi/4}|1_-\rangle,e^{+i\pi/4}|0_-\rangle\}\,, \tag{34}$$

presented in Fig. 4. The operations A and B correspond to braiding operations effected only on the + or the − topological qubits. These are single qubit operations. In contrast, if we effect braiding simultaneously in both qubits spaces we have two options. Either the direction of braiding is the same or opposite. These two possible topologically protected operations in the joint qubit space are described by the C and D configurations. The above set of protected operations do not suffice for performing universal TQC due to the Ising nature of the MFs [6, 50]. Nevertheless, the presence of the additional topological qubit on the *same* wire can be useful for performing braiding operations. So far, several methods for performing braiding have been proposed, including networks of topological wires [1, 51] where neighbouring MFs can be controllably coupled in order to perform a MF exchange. In the present case, the additional $\mathcal{O}_u$ protected MFs can constitute a reservoir of MFs that could reduce the number of complementary wires that one needs for performing adiabatic operations using these protocols. In addition, the presence of the extra pair of MFs can be also prominent for creating phase gate operations. A standard theoretical proposal [6] for implementing a phase gate for two separated MFs, prescribes to bring the MFs to a finite distance in order to let them hybridize into a finite energy fermionic state. Due to the time evolution of the finite energy state, a phase gate operation will be implemented on the topological qubit when the MFs reseparate. In the presence of a hidden symmetry $\mathcal{O}_u$, one does not have to change the distance of the MFs any more. By controllably switching off the hidden symmetry $\mathcal{O}_u$, one hybridizes the two MFs of the same edge, for instance $\gamma_{a\pm}$, so to end up with a single MF. Depending on the details of the hidden symmetry breaking and restoration procedures, one may retrieve a phase gate operation. Of course a detailed investigation of these possibilties is required.

The alternative TQC routes described above depend delicately and crucially on the robustness of this hidden symmetry. As we have already mentioned, it is desirable to find a system that has a hidden symmetry related to a degree of freedom such as a band index. For example, in the case of a two-band topological superconductor where only intraband matrix elements appear in the Hamiltonian, the system splits into two irreducible subsystems similarly to the situation described above. As a matter of fact, multiband systems such as the Fe-based high-Tc superconductors [52], offer a promising way out. The latter materials are supposed to exhibit intra-band superconductivity (usually a 4-band [53] or a 5-orbital [54] picture is adequate) and consequently we may obtain a number of disconnected sub-systems. If we manage to render each of these superconducting sub-systems topological, we will be in a position to apply the topological quantum computing protocols discussed in the previous paragraph. Recently, a proposal concerning topological superconductivity based on iron-based superconductors has been put forward [55]. However, in that work an iron-based superconductor was used to induce superconductivity by proximity effects on a Rashba-semiconductor. Instead, the situation that I envisage involves intrinsic multiband topological superconductivity in the iron pnictide superconductor itself.



FIG. 4: Allowed topologically protected operations for two topological qubits ± corresponding to the ±1 eigenvalues of a unitary hidden symmetry operator $\mathcal{O}_u$ satisfying $\mathcal{O}_u^2 = \mathrm{I}$. The standard braiding operations A and B describe single qubit operations. C and D correspond to operations in the joint qubit space. In operation C, the ± pairs of Majorana fermions are braided in the same direction while in case D in the opposite.

VII. CONCLUSION

I have performed a detailed analysis of the accessible topological superconducting phases that can occur from the combination of inhomogeneous Rashba spin-orbit coupling, magnetic order and superconductivity. By exploring the landscape of the possible topological phases I proposed new quasi-1d systems prominent for realizing MFs, based on Rashba spin-orbit coupling and $\mathcal{T}$ violating superconductivity, without the demand for any kind of magnetic order. Moreover, I pinpointed the significance of emergent unitary and anti-unitary hidden symmetries and revealed the topological implications that they lead to. Finally, I discussed alternative topological quantum computing pathways that open up in the presence of a unitary hidden symmetry and suggested that Fe-based multiband superconductors could be a potential candidate for these implementations.

Note

While the present manuscript was under completion, a preprint concerning the classification of topological insulators and superconductors in the presence of a reflection symmetry appeared [56].

Acknowledgements

I am grateful to Gerd Schön, Alexander Shnirman, Jens Michelsen and Daniel Mendler for the motivation and the great support. Furthemore, I am also indebted to Xiaoliang Qi, Ivar Martin, Alberto Morpurgo, Andreas Heimes, Lingzhen Guo, Juan Atalaya, Boris Narozhny, Mathias Scheurer, Bhilahari Jeevanesan and Elio König for numerous suggestions and enlightening discussions. In addition, I acknowledge financial support from the EU project NanoCTM.

[1] Fu Liang and Kane C L 2008 *Phys. Rev. Lett.* **100** 096407.
[2] Mourik V, Zuo K, Frolov S M, Plissard S R, Bakkers E P A M and Kouwenhoven L P 2012 *Science* **336** 6084; Rokhinson L P, Liu Xinyu, Furdyna J K 2012 *Nature Physics* **8** 795; Deng M T, Yu C L, Huang G Y, Larsson M, Caroff P, Xu H Q 2012 *Nano Lett.* **12** 6414; Das A, Ronen Y, Most Y, Oreg Y, Heiblum M and Shtrikman H 2012 arXiv:1205.7073.
[3] Hasan M Z and Kane C L 2010 *Rev. Mod. Phys.* **82** 3045; Qi X L and Zhang S C 2011 *Rev. Mod. Phys.* **83** 1057.
[4] Lutchyn R M, Sau J D and Das Sarma S 2010 *Phys. Rev. Lett.* **105** 077001.
[5] Oreg Y, Refael G and von Oppen F 2010 *Phys. Rev. Lett.* **105** 177002.
[6] Kitaev A Yu 2003 *Annals Phys.* **303** 2; Nayak C, Simon S H, Stern Ady, Freedman M and Das Sarma S 2008 *Rev. Mod. Phys.* **80** 1083; Ivanov D A 2001 *Phys. Rev. Lett.* **86** 268.
[7] Alicea J, Oreg Y, Refael G, von Oppen F and Fisher M P A 2011 *Nature Physics* **7** 412.
[8] Loss D and Di Vincenzo D P 1998 *Phys. Rev. A* **57** 120; Reilly D J, Taylor J M , Laird E A, Petta J R, Marcus C M, Hanson M P and Gossard A C 2008 *Phys. Rev. Lett.* **101** 236803.
[9] Makhlin Y, Schön G and Shnirman A 2001 *Rev. Mod. Phys.* **73** 357; Wellstood F C, Urbina C and Clarke J 1987 *Appl. Phys. Lett.* **50** 772.
[10] Kitaev A Yu 2001 *Phys. Usp.* **44**, 131.
[11] Fu Liang and Kane C L 2009 *Phys. Rev. B* **79** 161408; Jiang L, Pekker D, Alicea J, Refael G, Oreg Y and von Oppen F 2011 *Phys. Rev. Lett.* **107**, 236401.
[12] Kotetes P, Shnirman A and Schön G 2012 arXiv:1207.2691 (to appear in JKPS); Jiang L, Pekker D, Alicea J, Refael G, Oreg Y, Brataas A and von Oppen F 2012 *Phys. Rev. B* **87**, 075438; Meng Q, Shivamoggi V, Hughes T L, Gilbert M J and Vishveshwara S 2012 arXiv:1206.1295.
[13] San-Jose P, Prada E and Aguado R 2012 *Phys. Rev. Lett.* **108** 257001; Hützen R, Zazunov A, Braunecker B, Levy Yeyati A and Egger R 2012 *Phys. Rev. Lett.* **109** 166403; Ojanen T 2012 *Phys. Rev. Lett.* **109** 226804; Domínguez F, Hassler F and Platero G 2012 *Phys. Rev. B* **86** 140503(R); Mai S, Kandelaki E, Volkov A and Efetov K 2013 *Phys. Rev. B* **87** 024507; Pikulin D I and Nazarov Y V 2012 *Phys. Rev. B* **86** 140504(R); N. Didier, M. Gibertini, A. G. Moghaddam, J. König, and R. Fazio 2012 arXiv:1202.6357; Ojanen T 2012 arXiv:1210.3990.
[14] Leijnse M and Flensberg K 2011 *Phys. Rev. Lett.* **107** 210502.
[15] Lu Y M and Wang Z 2013 *Phys. Rev. Lett.* **110** 096403; Sela E, Altland A and Rosch A 2011 *Phys. Rev. B* **84**, 085114; Neupert T, Onoda S and Furusaki A 2010 *Phys. Rev. Lett.* **105** 206404; Chung S B, Zhang H J, Qi X L and Zhang S C 2011 *Phys. Rev. B* **84** 060510; Nersesyan A A and Tsvelik A M 2011 arXiv:1105.5835.
[16] Tsvelik A M 2011 arXiv:1106.2996v1.
[17] Choy T P, Edge J M, Akhmerov A R and Beenakker C W J 2011 *Phys. Rev. B* **84**, 195442.
[18] Kjaergaard M, Wölms K and Flensberg K 2012 *Phys. Rev. B* **85**, 020503.
[19] Martin I and Morpurgo A F 2012 *Phys. Rev. B* **85**, 144505.

[20] Nandkishore R and Levitov L 2012 *Phys. Rev. B* **82** 115124.
[21] Ramazashvili R 2008 *Phys. Rev. Lett.* **101** 137202.
[22] Majorana E 1937 *Nuovo Cimento* **5** 171.
[23] Wilczek F 2009 *Nature Physics* **5** 614.
[24] Read N and Green D 2000 *Phys. Rev. B* **61**, 10267.
[25] Michelsen J and Grein R 2012 arXiv:1208.1090; Sau J D, Lutchyn R M, Tewari S and Das Sarma S 2010 *Phys. Rev. B* **82** 094522; Stanescu T D, Sau J D, Lutchyn R M and Das Sarma S 2010 *Phys. Rev. B* **81** 241310(R); Grein R, Michelsen J and Eschrig M 2011 arXiv:1111.0445.
[26] Here, the characterization "unconventional" is assigned to order parameters of density wave or superconducting phases, that carry finite angular momentum such as p-wave, d-wave, etc.
[27] Nayak C 2000 *Phys. Rev. B* **62** 4880.
[28] Hsu C H, Raghu S and Chakravarty S 2011 *Phys. Rev. B* **84** 155111.
[29] Tewari S, Zhang C, Yakovenko V M and Das Sarma S 2008 *Phys. Rev. Lett.* **100** 217004.
[30] Kotetes P and Varelogiannis G 2010 *Phys. Rev. Lett.* **104** 106404; Zhang C, Tewari S, Yakovenko V M and Das Sarma S 2008 *Phys. Rev. B* **78** 174508.
[31] Kotetes P and Varelogiannis G 2008 *Phys. Rev. B* **78** 220509(R).
[32] Takei So, Fregoso B M, Galitski V and Das Sarma S 2013 *Phys. Rev. B* **87** 014504; Wong C L M and Law K T 2012 *Phys. Rev. B* **86** 184516.
[33] Altland A and Zirnbauer M R 1997 *Phys. Rev. B* **55** 1142.
[34] Ryu S, Schnyder A, Furusaki A and Ludwig A 2010 *New J. Phys.* **12** 065010.
[35] Volovik G E 2003 *Clarendon Press Oxford* "The Universe in a Helium Droplet".
[36] Qi X L, Hughes T and Zhang S C 2008 *Phys. Rev. B* **78** 195424.
[37] Fulde P and Ferrell R A 1964 *Phys. Rev.* **135** A550.
[38] Larkin A I and Ovchinnikov Y N 1965 *Sov. Phys. JETP* **20** 762; Agterberg D F and Tsunetsugu H 2008 *Nature Physics* **4** 639.
[39] Liu X J and Lobos A M 2013 *Phys. Rev. B* **87** 060504(R); Romito A, Alicea J, Refael G and von Oppen F *Phys. Rev. B* **85** 020502(R); Seradjeh B and Grosfeld E 2011 *Phys. Rev. B* **83** 174521.
[40] Potter C A and Lee P A 2010 *Phys. Rev. Lett.* **105** 227003; Potter C A and Lee P A 2011 *Phys. Rev. B* **83** 094525; Lutchyn R M, Stanescu T and Das Sarma S 2011 *Phys. Rev. Lett.* **106**; Tewari S, Stanescu T D, Sau J D and Das Sarma S 2012 *Phys. Rev. B* **86** 024504 (2012).
[41] Tewari S and Sau J D 2012 *Phys. Rev. Lett.* **109** 150408.
[42] Fang C, Gilbert M J and Bernevig B A 2012 *Phys. Rev. B* **86** 115112; Jadaun P, Xiao Di, Niu Q and Banerjee S K 2012 arXiv:1208.1472.
[43] Das T 2012 *Sci. Rep.* **2** 596.
[44] Kane C L and Mele E J 2005 *Phys. Rev. Lett.* **95** 146802; Kane C L and Mele E J 2005 *Phys. Rev. Lett.* **95** 226801; König M, Wiedmann S, Brüne C, Roth A, Buhmann H, Molenkamp L W, Qi X L and Zhang S C 2007 *Science* **318** 766; Roth A, Brüne C, Buhmann H, Molenkamp L W, Maciejko J, Qi X L and Zhang S C 2009 *Science* **325** 294.
[45] Kapitulnik A, Xia J, Schemm E and Palevski A 2009 *New J. Phys.* **11** 055060; Lutchyn R M, Nagornykh P and Yakovenko V M 2009 *Phys. Rev. B* **80** 104508; Goryo J 2008 *Phys. Rev. B* **78** 060501(R).
[46] Nandkishore R and Levitov L 2011 *Phys. Rev. Lett.* **107** 097402.
[47] Xiao Di, Chang M C and Niu Q 2010 *Rev. Mod. Phys.* **82** 1959.
[48] Goryo J and Maeda N 2011 *J. Phys. Soc. Jpn.* **80** 044707.
[49] Cheng M, Galitski V and Das Sarma S 2011 *Phys. Rev. B* **84** 104529; Goldstein G and Chamon C 2011 *Phys. Rev. B* **84** 205109; Budich J C, Walter S and Trauzettel B 2012 *Phys. Rev. B* **85** 121405; Rainis D and Loss D 2012 *Phys. Rev. B* **85** 174533; Schmidt M J, Rainis D and Loss D 2012 *Phys. Rev. B* **86** 085414.
[50] Bravyi S and Kitaev A 2005 *Phys. Rev. A* **71** 022316; Bravyi S 2006 *Phys. Rev. A* **73** 042313; Sau J D, Tewari S and Das Sarma S 2010 *Phys. Rev. A* **82** 052322.
[51] Sau J D, Clarke D J and Tewari S 2011 *Phys. Rev. B* **84** 094505; van Heck B, Akhmerov A R, Hassler F, Burrello M and Beenakker C W J 2012 *New J. Phys.* **14** 035019; Halperin B I, Oreg Y, Stern Ady, Refael G, Alicea J and von Oppen F 2012 *Phys. Rev. B* **85** 144501.
[52] Kamihara Y, Watanabe T, Hirano M and Hosono H 2008 *J. Am. Chem. Soc.* **130** 3296; D. J. Singh and M.H. Du, Phys. Rev. Lett. 100, 237003 (2008); Mazin I I and Schmalian J 2009 *Physica C* **469** 614.
[53] Korshunov M M and Eremin I 2008 *Phys. Rev. B* **78** 140509(R); Aperis A, Kotetes P, Varelogiannis G and Oppeneer P M 2011 *Physical Review B* **83** 092505.
[54] Kuroki K, Onari S, Arita R, Usui H, Tanaka Y, Kontani H and Aoki H 2008 *Phys. Rev. Lett.* **101** 087004.
[55] Zhang F, Kane C L and Mele E J 2012 arXiv:1212.4232.
[56] Chiu C K, Yao Hong and Ryu S 2013 arXiv:1303.1843.